Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



NSF® Certified Environmental Product Declaration www.nsf.org

Program Operator NSF® International
EPD Registration Number EPD 10834
Date of Publication 7/1/2023
Date of Validity 7/1/2028

This group of products include high durability upholstery products.

An EPD should provide current information, and may be updated if conditions change. The stated validity is therefore subject to the continued registration and publication at www.environdec.com.

Knoll Textiles

At KnollTextiles, we advocate for business practices that reduce the use, manufacture, and sale of products that have the potential to cause environmental damage or negatively impact human health and safety. We are dedicated to environmental stewardship and transparency, and as such, we prioritize lowemitting materials, recycled content, natural fibers, and the removal of Chemicals of Concern from our products.

Deepening our commitment to material, production, and operational transparency, we have engaged in Life Cycle Assessments (LCA) for selected products within our line. The LCA is a comprehensive quantitative analysis of the environmental impacts of the entire life cycle of our products. This information will be presented in product-specific Environmental Product Declarations (EPD) that report on the whole-life sustainability impact of operational, embodied, and long-term carbon.



This declaration is an environmental product declaration (EPD) in accordance with ISO 14025 and EN 15804 + A2.

Roul Textiles

According to ISO 14025 and EN 15804 + A2

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

EN 15804 +A2

EPDs rely on Life Cycle Assessment (LCA) to provide information on a number of environmental impacts of products over their life cycle. The EPD owner has the sole ownership, liability and responsibility of the EPD. Exclusions: EPDs do not indicate that any environmental or social performance benchmarks are met, and there may be impacts that they do not encompass. LCAs do not typically address the site-specific environmental impacts of raw material extraction, nor are they meant to assess human health toxicity. EPDs can complement but cannot replace tools and certifications that are designed to address these impacts and/or set performance thresholds – e.g., Type 1 certifications, health assessments and declarations, environmental impact assessments, etc. Accuracy of Results: EPDs regularly rely on estimations of impacts, and the level of accuracy in estimation of effect differs for any particular product line and reported impact. Comparability: EPDs are not comparative assertions and are either not comparable or have limited comparability when they cover different life cycle stages, are based on different product category rules or are missing relevant environmental impacts. EPDs from different programs may not be comparable. The ranges of the LCIA results are less than 10%.

EPD PROGRAM AND PROGRAM OPERATOR NAME, ADDRESS, LOGO, AND WEBSITE	NSF International, 789 N. Dixboro Ro	d, Ann Arbor, MI 4810	5,www.nsf.org					
PRODUCT CATEGORY RULES (PCR):	EN 15804 +A2							
INDEPENDENT THIRD-PARTY VERIFICATION OF THE DECLARATION AND DATA, ACCORDING TO ISO 14025:2006	EPD Process Certific	tification EPD verification						
DECLARATION NUMBER	EPD 10834							
DECLARED PRODUCT & DECLARED UNIT	Alignment, Alignment Backed, Beacc Messa, Plaidtastic, Prep, Quest, Ref Declared Unit = 1 m^2 of fabric (Drap	lect, Samos, Soiree,						
REFERENCE PCR AND VERSION NUMBER	EN 15804:2012+A2:2019: Sustainab for the product category of constructi		orks — Environmental Product Declarations — Core rules					
DESCRIPTION OF PRODUCT APPLICATION/USE	Knoll products are primarily used in	commercial and resid	ential settings.					
PRODUCT RSL DESCRIPTION	Not applicable for this product category	ory						
MARKETS OF APPLICABILITY	Global							
DATE OF ISSUE	July 1, 2023							
PERIOD OF VALIDITY	5 years							
EPD TYPE	Product Specific							
DATASET VARIABILITY	N/A							
EPD SCOPE	Cradle-to-Gate with Options							
YEAR(S) OF REPORTED PRIMARY DATA	2020							
LCA SOFTWARE & VERSION NUMBER	GaBi 10.6							
LCI DATABASE(S) & VERSION NUMBER	GaBi Sphera database, Service Pac	k 35						
LCIA METHODOLOGY & VERSION NUMBER	TRACI 2.1; CML 4.1							
PCR REVIEW WAS CONDUCTED BY:	EN15804+A2 Core PCR was develo	ped by the European	Committee for Standardization (CEN)					
This declaration was independently verified in accordance 15804 +A2 serves as the core PCR. INTERNAL This life cycle assessment was conducted in accordance PCR by:] EXTERNAL	Tony Favilla, NSF C	ertification, LLC Afavilla@nsf.org					
This life cycle assessment was independently verifie	d in accordance with ISO 14044 and							
the reference PCR by:	I I I I I I I I I I I I I I I I I I	Jack Geibig, EcoFo	m, LLC jgeibig@ecoform.com					

Environmental declarations from different programs (ISO 14025) may not be comparable.

Comparison of the environmental performance using EPD information shall consider all relevant information modules over the full life cycle of the products within the building

This PCR allows EPD comparability only when the same functional requirements between products are ensured and the requirements of EN 15804 are met. It should be noted that different LCA software and background LCI datasets may lead to differences results for upstream or downstream of the life cycle stages declared.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



Knoll Textiles

According to ISO 14025 and EN 15804 + A2

General Information

EN 15804 +A2

Description of Company/Organization

We at KnollTextiles use modern design to connect people to their work, their lives, their world. Since 1938, we have been recognized internationally for creating workplace and residential furnishings that inspire, evolve, and endure.

Today, our commitment to modern design, our understanding of the nature of work, and our dedication to sustainable design have yielded a unique portfolio of thoughtful products that respond and adapt to changing needs.

Product Description

KnollTextiles has many products which can be used in many different applications. This gives our customer many choices for the products functionality. The post-consumer recycled polyester fabrics covered in this EPD are the same across their product lines, where they differ is their final pattern. Due to their versatility, the products can be used for drapery, panelling, or for upholstery.

Manufacturer Specific EPD

This product-specific EPD was developed based on the cradle-to-grave with options (modules A1-A4 & C1-D) Life Cycle Assessment. The EPD accounts for raw material extraction and processing, transport, product manufacturing, distribution, and end-of-life. Manufacturing data were gathered directly from company personnel. When updated company-specific data were not available, a proxy was used. Product grouping was considered appropriate if the individual product(s) have a similar application and material composition. Any additional contracted facilities names have been withheld due to confidentiality.

Application

The Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II Drapery/Panel/Upholstery is typically used as a direct glue wallcovering, as well as unbacked for wrapped panel or indoor-outdoor upholstery use.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Material Composition

The primary product components and/or materials must be indicated as a percentage mass to enable the user of the EPD to understand the composition of the product in delivery status. There are no substances in these products listed in the Candidate List of Substances of Very High Concern for authorization. For more information see https://www.knoll.com/shop-textiles.

The average composition of the product(s) is as follows:

Material	Percentage	in mass (%)			
Post-Consumer Recycled Polyester	39.	00%			
Post-Industrial Recycled Polyester	32.	00%			
Polyester	29.00%				
Total	100.00%				
Total Mass	0.55	kg			

Technical Data

For the declared product, the following technical data in the delivery status must be provided with reference to the test standard:

	Technical Data
Composition	Post-Consumer Recycled Polyester
Weave	Dobby, Jacquard
Color Index	Various
Colorfastness Dry	Above ACT minimum Guidelines of Grade 4
Colorfastness Wet	Above ACT minimum Guidelines of Grade 3
ASTM E-84 Adhered	Meets ACT minimum Guidelines of Class A
ASTM E-84 Unadhered	Meets ACT minimum Guidelines of Class A
Lightfastness 40 hrs.	Meets ACT minimum Guidelines of Grade 4
ASTM C423 Acoustical	Please see individual product listing at https://www.knoll.com/shop-textiles for specific product rating
Clean Air	Tested to CDPH/BIFMA guidelines

Placing on the Market / Application Rules

These products can be used for Upholstery and/or Panel applications.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Methodological Framework

Declared Unit

The declaration refers to the Declared Unit of 1 m2 of Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II as specified in EN 15804 +A2. The declared unit was chosen to be consistent with the unit defined by EPD International PCR: Woven, Knitted or Crocheted Fabrics (of Synthetic Fibres) which was used to inform the EPD created under EN 15804 +A2 and because the use phase is out of scope.

Name	Value	Unit
Declared Unit	1 m2 (one se	quare meter) of fabric
Mass	0.55	kg

System Boundary

This is a cradle-to-gate with options Environmental Product Declaration. The following life cycle phases were considered:

Pro	duct Si	tage	Pro	truction ocess age		Use Stage						End of Life Stage*			Benefits and Loads Beyond the System Boundaries	
Raw material supply	Transport	Manufacturing	Transport from gate to the site	Construction/ installation process	Use	Maintenance Replacement Refurbishment Operational energy use Operational water use Transport				Transport	Waste processing	Disposal	Reuse-Recovery- Recycling potential			
A1	A2	А3	A4	A5	B1	B2	В3	B4	B5	B6	В7	C1	C2	С3	C4	D
X	Х	Х	Х	MND	MND	MND	MND	MND	MND	MND	MND	Х	Х	Х	Х	X

Description of the System Boundary Stages Corresponding to the PCR

(X = Included; MND = Module Not Declared)

Reference Service Life

The reference service life of a properly installed textile is not applicable because the use phase is out of scope. The building estimated service life is 75 years

Allocation

Allocation was determined on the declared unit for primary data, 1 m^2. For secondary data, cut-off methodology was used.

^{*}This includes provision of all materials, products and energy, packaging processing and its transport, as well as waste processing up to the end-of waste state or disposal of final residues.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

Certified Environmental Product Declaration Pr

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Cut-off Criteria

Processes whose total contribution to the final result, with respect to their mass and in relation to all considered impact categories, is less than 1% can be neglected. The sum of the neglected processes may not exceed 5% by mass of the considered impact categories.

For Hazardous Substances the following requirements apply:

- The Life Cycle Inventory (LCI) of hazardous substances will be included, if the inventory is available.
- If the LCI for a hazardous substance is not available, the substance will appear as an input in the LCI of the product, if its mass represents more than 0.1% of the product composition.
- If the LCI of a hazardous substance is approximated by modeling another substance, documentation will be provided.

This EPD is in compliance with the cut-off criteria. No known processes were neglected or excluded. Capital items for the production processes (machine, buildings, etc.) were not taken into consideration.

Data Sources

Primary data were collected for every process in the product system under the control of KnollTextiles. Secondary data from the GaBi Sphera database were utilized. These data were evaluated and have temporal, geographic, and technical coverage appropriate to the scope of the product category rule.

Data Quality

The data sources used are complete and representative of a global system boundary in terms of the geographic and technological coverage and are a recent vintage (i.e., less than ten years old). The data used for primary data are based on direct information sources of the manufacturer. Secondary data sets were used for raw materials extraction and processing, end of life, transportation, and energy production flows. Wherever secondary data is used, the study adopts critically reviewed data for consistency, precision, and reproducibility to limit uncertainty.

Period Under Review

The period under review is the full calendar year of 2020.

Treatment of Biogenic Carbon

The uptake and release of biogenic carbon throughout the product life cycle follows EN 15804. Biogenic carbon is only released through packaging materials

Comparability and Benchmarking

A comparison or an evaluation of EPD data is only possible if all data sets to be compared were created according to EN 15804 and the building context, respectively the product-specific characteristics of performance, are taken into account. Environmental declarations from different programs may not be comparable. Only EPDs that comply with EN 15804 are comparable. Full conformance with the PCR allows EPD comparability only when all stages of the fabric product's life cycle have been considered. However, variations and deviations are possible.

Units

The LCA results within this EPD are reported in SI units.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



Knoll Textiles

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2


Additional Environmental Information

Background data

For life cycle modeling of the considered products, the GaBi 10.6 Software System for Life Cycle Engineering, developed by Sphera, is used. The GaBi-database contains consistent and documented datasets which are documented in the online GaBi-documentation. To ensure comparability of results in the LCA, the basic data of GaBi database were used for energy, transportation and auxiliary materials.

Manufacturing

There are three basic steps required for fabric production. The first step is yarn production. This happens after the raw materials have been harvested and involves spinning the material into yarn and threads. For most manufacturing sites, this is completed using mechanized spinning wheels. After the raw materials are spun into yarn, the individual threads are joined together to form fabric. This step is called weaving and is done on a machine called a loom. After the fabric has been woven, it is removed from the loom and processed. In this step, the fabric is cleaned and treated with various chemicals and cleaners to remove impurities from the final product. The result of these steps is an all-white fabric. The last, optional, step is the dyeing of the fabric to its final color/pattern.



Certified Environmental Product Declaration www.nst.org

Knoll Textiles

According to ISO 14025 and EN 15804 + A2

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

EN 15804 +A2

Packaging

All packaging is fully recyclable. The packaging material is composed by cardboard, HDPE, LDPE, paper, and tape.

Material	Quantity (% By Weight)	Mass (g/m2)
Cardboard	93.32%	63.30
HDPE	0.34%	0.23
LDPE	6.10%	4.14
Paper	0.13%	0.09
Таре	0.10%	0.07
Total	100%	67.83

Transportation

Transport to Building Site (A4)		
Name	Value	Unit
Fuel type		Diesel
Liters of fuel	38	l/100km
Transport distance	800	km
Capacity utilization (including empty runs)	90	%
Gross density of products transported	-	kg/m³
Weight of products transported	-	kg
Volume of products transported	-	m ³
Capacity utilization volume factor	1.00	-

Disposal

End of life (C1-0	C4)	
Name	Value	Unit
Collected separately	0.11	kg
Collected as mixed construction waste	0.44	kg
Reuse	0.00	kg
Recycling	0.00	kg
Landfilling	0.44	kg
Incineration with energy recovery	0.11	kg
Energy conversion	44%	%
Material for final deposition	0.55	kg
Removals of biogenic carbon	0.00	kg

Re-use Phase

Re-Use, recovery, And/Or Recycling Potenti	al (D)	
Name	Value	Unit
Net energy benefit from energy recovery from waste treatment declared as exported energy in C3 (R>0.6)	0.00	MJ
Net energy benefit from thermal energy due to treatment of waste declared as exported energy in C4 (R<0.6)	0.00	MJ
Net energy benefit from material flow declared in C3 for energy recovery	0.00	MJ
Process and conversion efficiencies	-	_
Further assumptions for scenario development (e.g. further processing technologies, assumptions on correction factors);	-	-

According to ISO 14025 and EN 15804 + A2 **Knoll Textiles**

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

EN 15804 +A2

LCA Results per Declared Unit

				EN15804 +	· A2 Impact	Assessment					
Parameter	Parameter	Unit	A1	A2	A3	A4	C1	C2	C3	C4	D
	Total	kg CO2 eq	1.26E+02	4.38E-02	5.58E+02	9.96E-02	0.00E+00	5.21E-03	0.00E+00	3.66E-01	0.00E+00
	Fossil	kg CO2 eq	1.26E+02	4.38E-02	5.54E+02	9.96E-02	0.00E+00	5.21E-03	0.00E+00	8.70E-02	0.00E+00
Climate Change	Biogenic	kg CO2 eq	-4.65E-02	0.00E+00	4.50E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.79E-01	0.00E+00
	Land use and land use change	kg CO2 eq	8.76E-03	0.00E+00	1.31E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.92E-06	0.00E+00
Ozone Depletion		kg CFC-11 eq	5.82E-10	1.11E-12	7.59E-11	2.54E-12	0.00E+00	1.33E-13	0.00E+00	2.96E-14	0.00E+00
Acidification		Mole of H+ eq	9.99E-02	2.85E-04	2.13E-01	6.48E-04	0.00E+00	3.39E-05	0.00E+00	5.43E-04	0.00E+00
	Freshwater	kg P eq	1.48E-04	1.22E-08	1.33E-04	2.78E-08	0.00E+00	1.46E-09	0.00E+00	1.73E-05	0.00E+00
Eutrophication	Marine	kg N eq	3.52E-02	1.10E-04	2.07E-01	2.49E-04	0.00E+00	1.30E-05	0.00E+00	2.79E-04	0.00E+00
	Terrestrial	Mole of N eq	3.69E-01	1.20E-03	1.04E+00	2.72E-03	0.00E+00	1.42E-04	0.00E+00	2.37E-03	0.00E+00
Photochemica I ozone formation	Human Health	kg NMVOC eq	9.90E-02	3.23E-04	3.07E-01	7.34E-04	0.00E+00	3.84E-05	0.00E+00	3.06E-04	0.00E+00
Depletion of	Mineral and metals	kg Sb eq	7.70E-06	0.00E+00	3.24E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.37E-10	0.00E+00
abiotic resources	Fossils	MJ	2.25E+03	5.51E-01	8.33E+03	1.25E+00	0.00E+00	6.56E-02	0.00E+00	2.11E-01	0.00E+00
Water Use		m ³ world eq	2.33E+01	0.00E+00	-4.18E+01	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.43E-02	0.00E+00
Particulate matter		Disease incidences	1.16E-06	1.12E-09	2.72E-06	2.56E-09	0.00E+00	1.34E-10	0.00E+00	4.08E-09	0.00E+00
lonizing Radiation ¹	Human Health	kBq U235 eq.	1.93E+01	9.68E-21	5.12E-01	2.20E-20	0.00E+00	1.15E-21	0.00E+00	3.18E-04	0.00E+00
Ecotoxicity ²	Freshwater	CTUe	5.39E+02	2.32E+00	8.91E+02	5.28E+00	0.00E+00	2.76E-01	0.00E+00	4.08E+00	0.00E+00
lumana Tandaka	Cancer	CTUh	1.28E-08	1.16E-11	5.72E-08	2.64E-11	0.00E+00	1.38E-12	0.00E+00	4.58E-11	0.00E+00
Human Toxicity	Non-cancer	CTUh	5.97E-07	1.09E-09	1.39E-06	2.49E-09	0.00E+00	1.30E-10	0.00E+00	5.29E-09	0.00E+00
Land Use ²		Pt	7.41E+02	0.00E+00	3.18E+01	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.11E-02	0.00E+00

Disclaimer 1 - This impact category deals mainly with the eventual impact of low doze ionizing radiation on human health of the nuclear fuel cycle. It does not consider effects due to possible nuclear accidents, occupational exposure, nor due to radioactive waste disposal in underground facilities, Potential ionizing radiation from the soil, from radon, and from some construction materials is also not measured by this indicator.

Disclaimer 2 - The results of this environmental impact indicator shall be used with case as the uncertainties on these results are high, or as there is limited experienced with the indicator.

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

KnollTextiles

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Results shown below were calculated using TRACI 2.1 Methodology.

	TRACI 2.1 Impact Assessment													
Parameter	Parameter	Unit	A 1	A2	A3	A4	C1	C2	C3	C4	D			
	Fossil	kg CO ₂ -Eq.	1.28E+02	4.28E-02	5.36E+02	9.74E-02	0.00E+00	5.09E-03	0.00E+00	2.10E-01	0.00E+00			
OWE	Biogenic	kg CO ₂ -Eq.	7.80E-02	0.00E+00										
GWP	Land Use and Land Transformation	kg CO ₂ -Eq.	1.22E+02	4.18E-02	5.09E+02	9.51E-02	0.00E+00	4.98E-03	0.00E+00	2.24E-01	0.00E+00			
	Total	kg CO ₂ -Eq.	2.50E+02	8.46E-02	1.05E+03	1.93E-01	0.00E+00	1.01E-02	0.00E+00	4.34E-01	0.00E+00			
ODP	Depletion potential of the stratospheric ozone layer	kg CFC-11 Eq.	7.33E-12	1.62E-12	7.76E-11	3.68E-12	0.00E+00	1.93E-13	0.00E+00	7.36E-16	0.00E+00			
AP Air	Acidification potential for air emissions	kg SO ₂ -Eq.	1.16E-01	2.57E-04	2.79E-01	5.85E-04	0.00E+00	3.06E-05	0.00E+00	1.30E-03	0.00E+00			
EP	Eutrophication potential	kg N-Eq.	1.47E-02	1.42E-05	1.42E-01	3.24E-05	0.00E+00	1.70E-06	0.00E+00	5.04E-04	0.00E+00			
SP	Smog formation potential	kg O ₃ -Eq.	2.15E+00	7.08E-03	6.32E+00	1.61E-02	0.00E+00	8.42E-04	0.00E+00	4.65E-03	0.00E+00			
FFD	Fossil Fuel Depletion	MJ-surplus	1.58E+02	7.57E-02	1.20E+03	1.72E-01	0.00E+00	9.01E-03	0.00E+00	4.24E-02	0.00E+00			

^{*}All disposal stages have been considered and only those with non-zero values have been reported

Results shown below were calculated using CML Methodology.

	m below were calculated t		3,	CML 4.	1 Impact As	sessment					
Parameter	Parameter	Unit	A1	A2	А3	A4	C1	C2	C3	C4	D
GWP	Global warming potential	kg CO ₂ -Eq.	1.28E+02	4.29E-02	5.41E+02	9.76E-02	0.00E+00	5.11E-03	0.00E+00	2.25E-01	0.00E+00
ODP	Depletion potential of the stratospheric ozone layer	kg CFC-11 Eq.	3.72E-10	1.62E-12	8.88E-11	3.68E-12	0.00E+00	1.92E-13	0.00E+00	4.28E-14	0.00E+00
AP Air	Acidification potential for air emissions	kg SO₂-Eq.	1.04E-01	2.11E-04	1.68E-01	4.80E-04	0.00E+00	2.51E-05	0.00E+00	4.69E-04	0.00E+00
EP	Eutrophication potential	kg(PO ₄) ³ -Eq.	1.68E-02	3.76E-05	6.23E-02	8.56E-05	0.00E+00	4.48E-06	0.00E+00	5.62E-04	0.00E+00
POCP	Formation potential of tropospheric ozone photochemical oxidants	kg ethane-Eq.	1.13E-02	2.47E-05	3.77E-02	5.61E-05	0.00E+00	2.94E-06	0.00E+00	1.06E-04	0.00E+00
ADPE	Abiotic depletion potential for non-fossil resources	kg Sb-Eq.	2.73E-05	1.78E-11	1.90E-04	4.05E-11	0.00E+00	2.12E-12	0.00E+00	1.05E-08	0.00E+00
ADPF	Abiotic depletion potential for fossil resources	MJ	1.66E+03	5.46E-01	8.04E+03	1.24E+00	0.00E+00	6.50E-02	0.00E+00	3.35E-01	0.00E+00

^{*}All disposal stages have been considered and only those with non-zero values have been reported

Results below contain the resource use throughout the life cycle of the product.

Resource Use											
Parameter	Parameter	Unit	A1	A2	А3	A4	C1	C2	C3	C4	D
SM	Use of secondary material	kg	3.90E-04	0.00E+00							
RSF	Use of renewable secondary fuels	MJ	0.00E+00								
NRSF	Use of nonrenewable secondary fuels	MJ	0.00E+00								
FW	Use of net fresh water	m ³	6.12E-01	0.00E+00	9.07E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.62E-04	0.00E+00
PERE	Use of renewable primary energy	MJ	3.54E+02	0.00E+00	1.74E+01	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.30E-02	0.00E+00
PERM	Use of renewable primary energy resources used as raw materials	MJ	0.00E+00								
PERT	Total Use of Renewable primary energy	MJ	3.54E+02	0.00E+00	1.74E+01	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.30E-02	0.00E+00
PENRE	Use of Non-renewable Primary Energy	MJ	2.26E+03	5.51E-01	8.34E+03	1.25E+00	0.00E+00	6.56E-02	0.00E+00	2.13E-01	0.00E+00
PENRM	Use of non-renewable primary energy resources used as raw materials	MJ	0.00E+00								
PENRT	Total Use of Non- renewable Primary Energy	MJ	2.26E+03	5.51E-01	8.34E+03	1.25E+00	0.00E+00	6.56E-02	0.00E+00	2.13E-01	0.00E+00

^{*}All disposal stages have been considered and only those with non-zero values have been reported

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



Knoll Textiles

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Results below contain the output flows and wastes throughout the life cycle of the product.

Output Flows and Waste Categories											
Parameter	Parameter	Unit	A1	A2	A3	A4	C1	C2	C3	C4	D
HWD	Hazardous waste disposed	kg	1.20E-07	0.00E+00	6.08E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.43E-11	0.00E+00
NHWD	Non-hazardous waste disposed	kg	5.67E-01	0.00E+00	5.71E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.04E-01	0.00E+00
HLRW	High-level radioactive waste	kg or m ³	2.04E-01	0.00E+00	5.23E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.27E-06	0.00E+00
ILLRW	Intermediate- and low- level radioactive waste	kg or m ³	0.00E+00								
CRU	Components for re-use	kg	0.00E+00								
MR	Materials for recycling	kg	0.00E+00								
MER	Materials for energy recovery	kg	0.00E+00								
EE	Recovered energy exported from system	MJ	0.00E+00								

^{*}All disposal stages have been considered and only those with non-zero values have been reported

Results below contain direct greenhouse gas emissions and removals throughout the life cycle of the product.

Resource Use											
Parameter	Parameter	Unit	A 1	A2	А3	A4	C1	C2	C3	C4	D
BCRP	Biogenic Carbon Removal from Product	kg CO ₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
BCEP	Biogenic Carbon Emissions from Product	kg CO ₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
BCRK	Biogenic Carbon Removal from Packaging	kg CO ₂	7.80E-02	0.00E+00							
BCEK	Biogenic Carbon Emissions from Packaging	kg CO₂	0.00E+00	0.00E+00	7.80E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
BCEW	Biogenic Carbon Emissions from Combustion of Waste from Renewable Sources Used in Production Process	kg CO₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
CCE	Calcination Carbon Emissions	kg CO ₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
CCR	Carbonation Carbon Removal	kg CO ₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
CWNR	Carbon Emissions from Combustion of Waste from Non-renewable Sources Used in Production Process	kg CO₂	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00

^{*}All disposal stages have been considered and only those with non-zero values have been reported

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter

Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep,

Certified Environmental Product Declaration www.nst.org


KnollTextiles

According to ISO 14025 and EN 15804 + A2

Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II EN 15804 +A2

LCA Interpretation

The production life cycle stage (A3) dominates the impacts across all impact categories. This is due to the upstream production of fabrics and the fuel usage from the manufacturing facilities. There are no material impacts because the products are made of 100% recycled polyester and their impacts have already been accounted for in their previous life cycle.



Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

Exertified product Declaration wavestage

Knoll Textiles

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Additional Environmental Information

Environmental and Health During Manufacturing

There are no known health and safety concerns when manufacturing the fabrics in this group. There are no emissions to indoor air after or during installation

Environmental and Health During Installation

There is no known impact on health with these fabrics when being applied to furniture. Most of the products in this grouping are certified as Clean Air Gold for their low VOC emissions.

Extraordinary Effects

Fire

No danger to the environment can be anticipated.

Water

Contains no known substances that have any impact on water in case of flood.

Mechanical Destruction

No danger to the environment can be anticipated during mechanical destruction.

Delayed Emissions

Global warming potential is calculated using the TRACI 2.1 and CML 4.1 impact assessment methodologies. Delayed emissions are not considered.

Environmental Activities and Certifications

Globally recognized for pioneering modern fabrics using advanced materials and techniques, KnollTextiles is proud to offer a wide selection of sustainably designed products. Dedicated to environmental stewardship and transparency, we strive to meet our customers' needs for sustainability, healthier materials and third-party certification.

Further Information

KnollTextiles 1235 Water Street East Greenville, PA 18041

Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter

Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep,

Certified Environmental Product Declaration www.nst.org

KnollTextiles

According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

Contact Information

Study Commissioner



Knoll
Vanessa Thompson
Sustainability Coordinator
120 West Pumping Station Road, Suite A
Quakertown, PA 18951
vthompso@knoll.com
www.knoll.com

LCA Practitioner



Sustainable Solutions Corporation 155 Railroad Plaza, Suite 203 Royersford, PA 19468 USA (+1) 610 569-1047 info@sustainablesolutionscorporation.com www.sustainablesolutionscorporation.com

Alignment, Alignment Backed, Beacon, Bistro, Circle Line, Commuter Cloth, Criss Cross, Delite, Durand, Helios, Messa, Plaidtastic, Prep, Quest, Reflect, Samos, Soiree, Tabloid, and Woodland II



According to ISO 14025 and EN 15804 + A2

EN 15804 +A2

References

- PCR Part A EPD International (2017) General Programme Instructions for the International EPD® System. Version 3.0, dated 2017-12-11.
- PCR Part B Woven, Knitted Or Crocheted Fabrics (Of Synthetic Fibres), The International EPD System, Published November 2020.
- GaBi 10.6 thinkstep.one. GaBi Life Cycle Assessment version 10.6 (software).
- ISO 14025 ISO 14025:2011-10, Environmental labels and declarations Type III environmental declarations Principles and procedures.
- ISO 14040 ISO 14040:2009-11, Environmental management Life cycle assessment Principles and framework.
- ISO 14044 ISO 14044:2006-10, Environmental management Life cycle assessment Requirements and guidelines.
- EN 15804 + A2 EN 15804:2012+A2:2019: Sustainability of construction works Environmental Product Declarations Core rules for the product category of construction product
- NSF 2015 NSF International National Center for Sustainability Standards, General Program Instructions, 2015.
- Characterizatio IPCC. 2020. Climate Change 2020. The Physical Science Basis. Cambridge University Press. (http://www.ipcc.ch/report/ar5/wg1/). n Method
- Characterizatio Hauschild M.Z., & Wenzel H. Environmental Assessment of Products. Springer, US, Vol. 2, 1998.
 n Method
- Characterizatio Heijungs R., Guinée J.B., Huppes G., Lankreijer R.M., Udo de Haes H.A., Wegener Sleeswijk A. Environmental Life Cycle Assessment of Products: Guide and Backgrounds. CML. Leiden University, Leiden, 1992.
- Characterizatio Jenkin M.E., & Hayman G.D. Photochemical ozone creation potentials for oxygenated volatile organic compounds: sensitivity to n Method variations in kinetic and mechanistic parameters. Atmospheric Environment. 1999, 33 (8) pp. 1275-1293.
- Characterizatio WMO. 1999. Scientific Assessment of Ozone Depletion: 1998, World Meteorological Organization Global Ozone Research and n Method Monitoring Project Report No. 44, WMO, Geneva.
- Characterizatio Standard Method for the Testing and Evaluation of Volatile Organic Chemical Emissions from Indoor Sources using Environmental n Method Chambers- version 1.2, January 2017.
- Characterizatio Intertek Clean Air Gold Certification, ISO 17065 accredited n Method